

Higher Spin Lifshitz Holography

Michael Gary

Institute for Theoretical Physics, TU Wien

Prague, 28.03.2014

Forthcoming Work, MG, D Grumiller, S Prohazka, S J Rey
JHEP 1211 (2012) 099 [arXiv:1209.2860] H Afshar, MG,
D Grumiller, R Rashkov, M Riegler

Outline

- 1 Motivation
- 2 Pure Gravity in AdS_3
- 3 Higher Spin Gravity in AdS_3
- 4 More General Backgrounds
- 5 Asymptotic Lifshitz
- 6 Conclusions

Motivation

- Higher Spin Holography with Lifshitz Scaling
- Role of Geometry in Higher Spin Theories
- Understanding the Mechanisms of Holography

3D Gravity

- Einstein-Hilbert action with negative cosmological constant

$$S = \frac{1}{16\pi G_N} \int_{\mathcal{M}} \sqrt{-g} \left(R + \frac{2}{\ell^2} \right)$$

- Vacuum solution is AdS_3

$$ds_{\text{AdS}}^2 = \ell^2 \left(-\cosh^2 \rho dt^2 + d\rho^2 + \sinh^2 \rho d\phi^2 \right)$$

- All solutions are locally AdS_3
- BTZ Black Hole

$$ds_{\text{BTZ}}^2 = \ell^2 \left[d\rho^2 - \left(e^{2\rho} + 16(M^2 - J^2) \right) dx^+ dx^- \right. \\ \left. + 4(M - J)(dx^+)^2 + 4(M + J)(dx^-)^2 \right]$$

Chern-Simons Formulation

- Gravity in Asymptotically AdS_3 can be formulated as $\mathfrak{sl}_2(\mathbb{R}) \oplus \mathfrak{sl}_2(\mathbb{R})$ Chern-Simons theory with level $k = \frac{1}{4G_N}$

$$S = \frac{k}{4\pi} (S_{\text{CS}}[A] - S_{\text{CS}}[\bar{A}])$$

$$S_{\text{CS}}[A] = \int_{\mathcal{M}} \text{tr} \left(A \wedge dA - \frac{2}{3} A^3 \right)$$

where

$$e = \frac{\ell}{2} (A - \bar{A}) \qquad \omega = \frac{1}{2} (A + \bar{A})$$

- Gauge transformations $\delta_\epsilon A = d\epsilon + [\epsilon, A]$, $\delta_{\bar{\epsilon}} \bar{A} = d\bar{\epsilon} + [\bar{\epsilon}, \bar{A}]$
- Diffeomorphisms generated by ξ^μ are given by

$$\epsilon = \xi^\mu A_\mu \qquad \bar{\epsilon} = \xi^\mu \bar{A}_\mu$$

Brown-Henneaux Boundary Conditions

- Denote \mathfrak{sl}_2 generators by $L_0, L_{\pm 1}$
- Convenient to partially gauge fix

$$A = g^{-1}dg + g^{-1}ag \quad \bar{A} = gdg^{-1} + g\bar{a}g^{-1} \quad g = e^{\rho L_0}$$

- Impose Asymptotic AdS boundary conditions

$$a = (L_1 + \mathcal{L}(x^+)L_{-1}) dx^+ + o(1)$$

$$\bar{a} = (L_{-1} + \bar{\mathcal{L}}(x^-)L_1) dx^- + o(1)$$

- Solutions include AdS, BTZ black holes, more

$$ds^2 = \ell^2 [d\rho^2 - (e^{2\rho} + e^{-2\rho} \mathcal{L}\bar{\mathcal{L}}) dx^+ dx^- + \mathcal{L}(dx^+)^2 + \bar{\mathcal{L}}(dx^-)^2 + \dots]$$

Canonical Analysis

- Locally, all solutions are flat, so gauge equivalent to the vacuum
- At the asymptotic boundary, some first class constraints become second class, and thus generate new states, rather than gauge transformations
- Asymptotic Symmetry Algebra is two copies of Virasoro with $c_L = c_R = 6k$

$$[L_n, L_m] = (n - m)L_{n+m} + \frac{c}{12}n(n^2 - 1)\delta_{n,-m}$$

- CFT vacuum defined by $L_n |0\rangle = 0$ for all $n \geq -1$ (similar for barred sector)
- States generated by $L_{n_1} \cdots L_{n_m} |0\rangle$ for $n_i < -1$, called boundary gravitons

Symmetries & Vacuum

- AdS solution preserves $3+3$ symmetries corresponding to the wedge algebra of the ASA, $\mathfrak{sl}_2 \oplus \mathfrak{sl}_2$, and is thus identified with the CFT vacuum
- All solutions locally preserve $3+3$ symmetries, since all solutions locally flat, but global realization is such that they excite infinite numbers of charges, not the wedge algebra

Higher Spin Generalization

- Enlarge \mathfrak{sl}_2 to \mathfrak{sl}_N
- Choice of embedding $\mathfrak{sl}_2 \hookrightarrow \mathfrak{sl}_N$ determines other field content
- Spins of other fields given by weight under gravitational \mathfrak{sl}_2 action
- Typical choice: Principal embedding, integer spins $2, \dots, N$.

$$\begin{aligned}g_{\mu\nu} &= \frac{1}{2} \text{tr} [e_\mu e_\nu] \\ \phi_{\mu\nu\rho} &= \text{tr} [e_{(\mu} e_\nu e_{\rho)}] \\ &\vdots\end{aligned}$$

\mathfrak{sl}_3 Conventions

- \mathfrak{sl}_2 generators $L_0, L_{\pm 1}$
- Spin 3 generators $W_0, W_{\pm 1}, W_{\pm 2}$
- Commutators

$$[L_n, L_m] = (n - m)L_{n+m} \quad [L_n, W_m] = (2n - m)W_{n+m}$$

$$[W_n, W_m] \propto L_{n+m}$$

- Traces

$$\text{tr}(L_n L_m) \propto \delta_{n, -m} \quad \text{tr}(W_n W_m) \propto \delta_{n, -m}$$

$$\text{tr}(L_n W_m) = 0$$

AdS Boundary Conditions

$$A = g^{-1}dg + g^{-1}ag \quad \bar{A} = gdg^{-1} + g\bar{a}g^{-1} \quad g = e^{\rho L_0}$$

$$a = (L_1 + \mathcal{L}(x^+)L_{-1} + \mathcal{W}(x^+)W_{-2}) dx^+ + o(1)$$

$$\bar{a} = (L_{-1} + \bar{\mathcal{L}}(x^-)L_1 + \bar{\mathcal{W}}(x^-)W_2) dx^- + o(1)$$

- Asymptotic Symmetry Algebra: two copies of \mathcal{W}_3 with central charges $c_L = c_R = 6k$
- Vacuum: metric is AdS_3 , spin-3 field is 0, invariant under $\mathfrak{sl}_3 \times \mathfrak{sl}_3$ symmetry

General Procedure

- Add boundary term to cancel variation of the action

$$S_{\text{CT}} = -\frac{k}{4\pi} \int_{\partial\mathcal{M}} \text{tr} \left(A^2 - \overline{A}^2 \right)$$

- Split connection into background and fluctuations
- Impose consistent boundary conditions on fluctuations. In particular
 - Find closed set of boundary condition preserving gauge transformations
 - Require finite, conserved, integrable asymptotic charges
- Determine Asymptotic Symmetry Algebra by computing Poisson Brackets and quantizing

Examples

- Lobachevsky boundary conditions

$$\hat{ds}^2 = d\rho^2 \pm dt^2 + \sinh^2 \rho d\phi^2$$

$\mathcal{W}_N^{(2)} \times \hat{\mathfrak{u}}(1)$ Asymptotic Symmetry Algebra

[1209.2860], [1211.4454]

- Null-Warped AdS₃ boundary conditions

$$\hat{ds}^2 = d\rho^2 + e^{2\rho} dt d\phi + \frac{9}{4} e^{4\rho} d\phi^2$$

$\mathcal{W}_3^{(2)}$ Asymptotic Symmetry Algebra

(Unpublished, with E Perlmutter and D Grumiller)

Lifshitz Geometry

- Lifshitz geometries are dual to Lifshitz field theories, which feature anisotropic scaling between space and time with a relative factor z

$$t \rightarrow \lambda^z t \qquad x \rightarrow \lambda x$$

- Metric

$$\begin{aligned} ds_z^2 &= \ell^2 \left(-r^{2z} dt^2 + \frac{dr^2}{r^2} + r^2 dx^2 \right) \\ &= \ell^2 \left(-e^{2z\rho} dt^2 + d\rho^2 + e^{2\rho} dx^2 \right) \end{aligned}$$

- Isometries and Lifshitz Algebra

$$\xi_{\mathbb{H}} = \partial_t \qquad \xi_{\mathbb{P}} = \partial_x \qquad \xi_{\mathbb{D}} = -zt\partial_t + \partial_\rho - x\partial_x$$

$$[\xi_{\mathbb{H}}, \xi_{\mathbb{P}}] = 0 \qquad [\xi_{\mathbb{D}}, \xi_{\mathbb{H}}] = z\xi_{\mathbb{H}} \qquad [\xi_{\mathbb{D}}, \xi_{\mathbb{P}}] = \xi_{\mathbb{P}}$$

$z = 2$ Lifshitz Background

- Background connection

$$\hat{a} = L_1 dx + \frac{4}{9} W_2 dt$$

$$\hat{\hat{a}} = L_{-1} dx + W_{-2} dt$$

- Background metric

$$ds^2 = \ell^2 \left(-e^{4\rho} dt^2 + d\rho^2 + e^{2\rho} dx^2 \right)$$

- Non-trivial background spin-3 field

$$\phi_{\mu\nu\lambda} dx^\mu dx^\nu dx^\lambda = -\frac{5\ell^3}{4} e^{4\rho} dt (dx)^2$$

Higher Spin Fluctuations on Lifshitz Background

- Boundary conditions

$$a^{(0)} = \left(4t\mathcal{W}L_0 - \mathcal{L}L_{-1} - \frac{16t^2}{9}\mathcal{W}W_2 + \frac{16t}{9}\mathcal{L}W_1 + \mathcal{W}W_{-2} \right) dx$$

$$\bar{a}^{(0)} = \left(-\bar{\mathcal{L}}L_1 - 9t\bar{\mathcal{W}}L_0 + \bar{\mathcal{W}}W_2 + 4t\bar{\mathcal{L}}W_{-1} - 9t^2\bar{\mathcal{W}}W_{-2} \right) dx$$

- Theory includes states with metrics that would not typically be called asymptotically Lifshitz
- On-shell solutions have explicit t -dependence

Asymptotic Symmetry Algebra

- Asymptotic charges are $\mathcal{L}(x), \mathcal{W}(x), \overline{\mathcal{L}}(x), \overline{\mathcal{W}}(x)$, t -independent
- Asymptotic Symmetry Algebra: two copies of \mathcal{W}_3 with central charges $c_L = c_R = 12k \text{tr}(L_0)^2 = \frac{3\ell}{2G_N}$

$$\begin{aligned}
 \delta_{\epsilon_L} \mathcal{L} &= \mathcal{L}' \epsilon_L + 2\mathcal{L} \epsilon'_L - \frac{k}{\pi} \epsilon_L^{(3)} \\
 \delta_{\epsilon_L} \mathcal{W} &= \mathcal{W}' \epsilon_L + 3\mathcal{W} \epsilon'_L \\
 \delta_{\epsilon_W} \mathcal{L} &= 2\mathcal{W}' \epsilon_W + 3\mathcal{W} \epsilon'_W \\
 \delta_{\epsilon_W} \mathcal{W} &= \left(\frac{3\pi}{k} \mathcal{L} \mathcal{L}' - \frac{3}{8} \mathcal{L}^{(3)} \right) \epsilon_W + \left(\frac{3\pi}{k} \mathcal{L} \mathcal{L}' - \frac{3}{8} \mathcal{L}^{(2)} \right) \epsilon'_W \\
 &\quad - \frac{45}{16} \mathcal{L}' \epsilon_W'' - \frac{15}{8} \mathcal{L} \epsilon_W^{(3)} + \frac{3k}{16\pi} \epsilon_W^{(5)}
 \end{aligned}$$

Symmetries of the Background

- Background is invariant under $8 + 8$ linearly independent gauge transformations of the form

$$\epsilon_L = l_{+1} - x l_0 + x^2 l_{-1}$$

$$\epsilon_W = w_{+2} - x w_{+1} + x^2 w_0 - x^3 w_{-1} + x^4 w_{-2}$$

$$\epsilon_{\bar{L}} = \bar{l}_{-1} - x \bar{l}_0 + x^2 \bar{l}_{+1}$$

$$\epsilon_{\bar{W}} = \bar{w}_{-2} - x \bar{w}_{-1} + x^2 \bar{w}_0 - x^3 \bar{w}_{+1} + x^4 \bar{w}_{+2}$$

- Special case: Lifshitz isometries

$$\xi_{\mathbb{H}} : \quad w_{+2} = \frac{4}{9} \quad \bar{w}_{-2} = 1$$

$$\xi_{\mathbb{P}} : \quad l_{+1} = 1 \quad \bar{l}_{-1} = 1$$

$$\xi_{\mathbb{D}} : \quad l_0 = 1 \quad \bar{l}_0 = 1$$

Symmetries & Global Structure

- Symmetries of the background enhanced to the full wedge algebra $\mathfrak{sl}_3 \times \mathfrak{sl}_3$, thus background is dual to the CFT vacuum on the plane
- All states locally have $8 + 8$ symmetries, but globally realized non-polynomially, leading to infinite towers of non-trivial charges
- No other states are invariant under precisely the complete wedge algebra

Interpretations and Duality

- Perturbative spectrum is equivalent to that of spin-3 gravity in AdS_3 : duality?
- Unclear if duality holds also at non-perturbative level
- Polynomial x and t dependence of boundary conditions and boundary condition preserving gauge transformations imply boundary should be non-compact

Conclusions

- Also looser boundary conditions for Lifshitz background, possibly related to $\mathcal{W}_3^{(2)}$
- Are there other backgrounds which are also related by duality?
- Do such dualities exist for more general higher spin theories?
- Metric and higher spin fields need to be placed on equal footing—all massless degrees of freedom
- To really talk about geometry, we should use a local probe (e.g. scalar field in $HS(\lambda)$ theory)

Thank You