

G. Zoupanos

NTUA Athens

# Finite Unified Theories

Predictions for the lightest Higgs mass

and other collider observables

SM very successful!

BUT with  $> 20$  free parameters

ad hoc Higgs sector



ad hoc Yukawa couplings

Best candidate for Physics? and SM

MSSM with  $> 100!$  free parameters

mostly in its SSB sector.

- cures problem of quadratic divergencies of the SM (hierarchy problem)
- restricts the Higgs sector



- SM with two - Higgs doublets

$$V = m_1^2 |H_1|^2 + m_2^2 |H_2|^2 + (m_3^2 H_1 H_2 + \text{h.c.})$$

$$+ \frac{1}{2} \lambda_1 (H_1^\dagger H_1)^2 + \frac{1}{2} \lambda_2 (H_2^\dagger H_2)^2$$

$$+ \lambda_3 (H_1^\dagger H_1) (H_2^\dagger H_2) + \lambda_4 (H_1 H_2) (H_1^\dagger H_2^\dagger)$$

$$+ \left\{ \frac{1}{2} \lambda_5 (H_1 H_2)^2 + [\lambda_6 (H_1^\dagger H_1) + \lambda_7 (H_1^\dagger H_2^\dagger)] (H_1 H_2) + \text{h.c.} \right\}$$

Supersymmetry provides tree level relations among couplings

$$\lambda_1 = \lambda_2 = \frac{1}{4} (g^2 + g'^2)$$

$$\lambda_3 = \frac{1}{4} (g^2 - g'^2), \quad \lambda_4 = -\frac{1}{4} g^2$$

$$\lambda_5 = \lambda_6 = \lambda_7 = 0$$

With  $v_1 = \langle \text{Re } H_1^0 \rangle$ ,  $v_2 = \langle \text{Re } H_2^0 \rangle$

$$\text{and } v_1^2 + v_2^2 = (246 \text{ GeV})^2, \quad \frac{v_2}{v_1} \equiv \tan \beta$$

$$\Rightarrow h^0, H^0, H^\pm, A^0$$

At tree level

$$M_{h, H^0}^2 = \frac{1}{2} \left\{ M_A^2 + M_Z^2 \mp \left[ (M_A^2 + M_Z^2)^2 - 4 M_A^2 M_Z^2 \cos^2 2\beta \right]^{1/2} \right\}$$

$$M_{H^\pm}^2 = M_W^2 + M_A^2$$

$$\Rightarrow \begin{cases} M_{h^0} < M_Z | \cos 2\beta | \\ M_{H^0} > M_Z \\ M_{H^\pm} > M_W \end{cases}$$

Radiative corrections

$$M_{h^0}^2 \simeq M_Z^2 \cos^2 2\beta + \frac{3g^2 m_t^4}{16\pi^2 M_W^2} \log \frac{m_{t1}^2 m_{t2}^2}{m_t^4}$$

- Finite Unified Theories  
(from Quantum Reduction  
of Couplings)
- Higher Dimensional Unified Theories  
and Coset Space Dimensional  
Reduction (Classical Reduction  
of Couplings)
- Fuzzy Extra Dimensions  
and Renormalisable Unified Theories

# Quantum

## Reduction of Couplings

Consider a GUT with

$g$  - gauge coupling

$g_i$  - other couplings (Yukawas, self-couplings)

Any relation among the couplings

$$\Phi(g, g_1, \dots) = \text{const}$$

which is RGI should satisfy

$$\frac{d}{dt} \Phi = 0, \quad t = \ln \mu$$

$$\frac{d}{dt} \Phi = \frac{\partial \Phi}{\partial g} \frac{dg}{dt} + \sum_i \frac{\partial \Phi}{\partial g_i} \frac{dg_i}{dt} = 0$$

which is equivalent to

$$\frac{dg}{b_g} = \frac{dg_1}{b_1} = \frac{dg_2}{b_2} = \dots \quad \begin{array}{l} \text{characteristic} \\ \text{system} \end{array}$$

$$\Rightarrow b_g \frac{d g_i}{d g} = b_i$$

Reduction  
egs  
Dehne  
Zimmermann

Demand power series solution to the REs

$$g_i = \sum_{n=0}^{\infty} \rho_i^{(n+1)} g^{2n+1}$$

Remarkably, uniqueness of these solutions can be decided already at 1-loop!

Assume

$$b_i = \frac{1}{16\pi^2} \left[ \sum_{j,k,l} b_i^{(1)jkl} g_j g_k g_l + \sum_{j \neq g} b_i^{(1)j} g_j g^2 \right] + \dots$$

$$b_g = \frac{1}{16\pi^2} b_g^{(1)} g^3 + \dots$$

Assume  $\rho_i^{(n)}$ ,  $n \leq r$  have been uniquely determined

To obtain  $\rho_i^{(r+1)}$ , insert  $g_i$  in REs and collect terms of  $O(g^{2r+1})$

$$\rightarrow \sum_{l \neq g} M(r)_i^l p_l^{(r+1)} = \text{lower order quantities known by assumption}$$

where

$$M(r)_i^l = 3 \sum_{j, k \neq g} b_i^{(1)jkl} p_j^{(1)} p_k^{(1)} + b_i^{(1)l} - (2r+1) b_g^{(1)l} \delta_i^l$$

$$0 = \sum_{j, k, l \neq g} b_i^{(1)jkl} p_j^{(1)} p_k^{(1)} p_l^{(1)} + \sum_{l \neq g} b_i^{(1)l} p_l^{(1)} - b_g^{(1)} p_i^{(1)}$$

$\Rightarrow$  for a given set of  $p_i^{(1)}$ , the  $p_i^{(n)}$  for all  $n > 1$  can be uniquely determined if

$$\det M(n)_i^l \neq 0$$

for all  $n$



Consider an  $SU(N)$  (non-susy) theory with

$\phi^i(N)$ ,  $\hat{\phi}_i(\bar{N})$  - complex scalars

$\psi^i(N)$ ,  $\hat{\psi}_i(\bar{N})$  - Weyl spinors

$T^a$  ( $a=1, \dots, N^2-1$ ) - "

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu} + i\sqrt{2} [g_Y \bar{\psi} T^a \psi - \hat{g}_Y \hat{\psi} T^a \hat{\phi}] + \text{h.c.} - V(\phi, \hat{\phi}),$$

$$V(\phi, \hat{\phi}) = \frac{1}{4} \lambda_1 (\phi^i \phi_i^*)^2 + \frac{1}{4} \lambda_2 (\hat{\phi}_i \hat{\phi}^{*i})^2 \\ + \lambda_3 (\phi^i \phi_i^*) (\hat{\phi}_j \hat{\phi}^{*j}) \\ + \lambda_4 (\phi^i \phi_j^*) (\hat{\phi}_i \hat{\phi}^{*j})$$

Searching for power series solution of the R.E.s we find

$$g_Y = \hat{g}_Y = g; \lambda_1 = \lambda_2 = \frac{N-1}{N} g^2; \lambda_3 = \frac{1}{2N} g^2; \lambda_4 = -\frac{1}{2} g^2 \\ \text{i.e. SUSY}$$

## $N=1$ gauge theories

Consider a chiral, anomaly free,  $N=1$  globally supersymmetric gauge th. based on a group  $G$  with gauge coupling  $g$ .

### Superpotential

$$W = \frac{1}{2} m_{ij} \phi^i \phi^j + \frac{1}{6} C_{ijk} \phi^i \phi^j \phi^k$$

$m_{ij}, C_{ijk}$  - gauge invariant tensors

$\phi^i$  - matter fields transforming as an

ir. rep.  $R_i$  of  $G$ .

Renormalization constants associated with  $W$

$$\phi^{oi} = (Z_j^i)^{1/2} \phi^j, \quad m_{ij}^o = Z_{ij}^{i's'} m_{i's'}^o, \quad C_{ijk}^o = Z_{ijk}^{i's'k'} C_{i's'k'}^o$$

$N=1$  non-renormalization thm ensures absence of mass and cubic-int-term infinities

$$Z_{i's'k'}^{ijk} Z_{i''}^{1/2 i'} Z_{j''}^{1/2 j'} Z_{k''}^{1/2 k'} = \delta_{(i''}^i \delta_{j''}^j \delta_{k''}^k)$$

$$Z_{i's'}^{ij} Z_{i''}^{1/2 i'} Z_{j''}^{1/2 j'} = \delta_{(i''}^i \delta_{j''}^j)$$

(In the background field method)

$$Z_g Z_v^{1/2} = 1$$

$\rightarrow$  Only surviving infinities are  $Z_{jj}^i(Z_v)$   
i.e. one infinity for each field.

The 1-loop  $\beta$ -function of the gauge coupling is

$$\beta_g^{(1)} = \frac{dg}{dt} = \frac{g^3}{16\pi^2} \left[ \sum_i \ell(R_i) - 3C_2(G) \right]$$

$\ell(R_i)$  - Dynkin index of  $R_i$

$C_2(G)$  - quadratic Casimir of the adjoint rep.

$\beta$ -functions of  $C_{ijk}$ . In order of the renormalization they are written with the anomalous dim. matrix  $\gamma_i^j$  of  $\phi^i$

$$\beta_{ijk}^{(1)} = \frac{dC_{ijk}}{dt} = C_{ije} \gamma_k^e + C_{ike} \gamma_j^e + C_{jke} \gamma_i^e$$

$$\gamma_i^j = Z^{-\frac{1}{2}k} \frac{d}{dt} Z^{\frac{1}{2}j}$$

$$= \frac{1}{32\pi^2} \left[ C^{jke} C_{ike} - 2g^2 C_2(R_i) \delta_i^j \right]$$

$C_2(R_i)$  - quadratic Casimir of  $R_i$

$$C^{ijk} = C^*_{ijk}$$

$$b_g^{(2)} = \frac{1}{(16\pi^2)^2} 2 g^5 \left[ \sum_i l(R_i) - 3 C_2(G) \right]$$

$$- \frac{1}{(16\pi^2)^2} \frac{g^3}{r} C_2(R_i) \left[ C^{jkl} C_{ikl} - 2g^2 C_2(R_i) \delta_{ij} \right]$$

$r: \text{tr} \sigma^{ab}$

Parke, West, Jones  
 Mezincescu, Yau  
 Machacek, Vaughn

$$\gamma^{(2)ij} = \frac{1}{(16\pi^2)^2} 2 g^4 C_2(R_i) \left[ \sum_i l(R_i) - 3 C_2(G) \right]$$

$$- \frac{1}{(16\pi^2)^2} \frac{1}{2} \left[ C^{ikl} C_{jklm} + 2g^2 (R^a)_m^i (R^a)_j^l \right]$$

$$\cdot \left[ C^{mpq} C_{lpq} - 2 \delta_l^m g^2 C_2(R_i) \right]$$

$$b_g^{NSVZ} = \frac{g^3}{16\pi^2} \left[ \frac{\sum_i l(R_i) (1 - 2\gamma_i) - 3 C_2(G)}{1 - g^2 C_2(G) / 8\pi^2} \right]$$

Novikov - Shifman - Vainshtein - Zakharov

# Finite Unification

Old days...

... divergences are "hidden under the carpet" (Dirac, Lects on Q.F.T., '64)

Recent years ...

... divergences reflect existence of a higher scale where new degrees of freedom are excited.

Not just artifacts of pert. th.

However the presence of quadratic divergences means that physics at one scale are very sensitive to unknown physics at higher scales.

→ SUSY ths which are free of quadratic divergences in spite of any experimental evidence...

→ Natural to expect that beyond unification scale the theory should be completely finite.

- $N=4$  → finite to all orders in pert.
- $N=2$  → only 1-loop contributions to  $\beta$ -function. Possible to arrange the spectrum so that theory is finite.

Multiplicities for massless irreducible reps with maximal helicity 1

$N$ $S_{\text{pin}}$	1	1	2	2	4
1	—	1	—	1	1
1/2	1	1	2	2	4
0	2	—	4	2	6

$$N=2 : b(g) = \frac{2g^3}{(4\pi)^2} \left( \sum_i T(R_i) - C_2(G) \right)$$

e.g.  $SU(N)$  with  $2N$  fundamental  
reps  $\rightarrow b(g) = 0$

$$SU(5) : p(5 + \bar{5}) ; q(10 + \bar{10}) ; r(15 + \bar{15})$$

with  $p + 3q + 7r = 10$

$$SO(10) : p(10 + \bar{10}) ; q(16 + \bar{16})$$

with  $p + 2q = 8$

$$E_6 : 4(27 + \bar{27})$$



# Finite Unified Theories

$$N=1$$

- 1-loop finiteness conditions

$$b_g^{(1)} = 0$$

$$\gamma_j^{(1)i} = 0 \text{ - anomalous dimensions of all chiral superfields}$$

- Exists complete classification of all chiral  $N=1$  models with  $b_g^{(1)} = 0$   
Hamidi - Patera - Schwarz  
Jiang - Zhou

- 1-loop finiteness Parkes-West  
Jones  
→ 2-loop finiteness Mezincescu

... Exist simple criteria  
 that guarantee all  
 loop finiteness  
 (vanishing of all-loop  
 beta functions)

Lucchesi-Piquet  
 Sibold

Ermushev  
 Kazakov  
 Tarasov

Leigh-Strassler

• All-loop finiteness (SUSY)  
 $\Rightarrow$  top mass  $\neq 0$  ✓

Kapetomakis  
 Mondragon  
 2  
 '92

---

 SUSY sector

• 1-loop finiteness conds

Jones  
 Meziurescu  
 Yao

(require in particular  
 universal soft SUSY  
 scalar masses)

$$(m^2)_j^i = \frac{1}{3} M M^* \delta_j^i$$

•• 1-loop finiteness

Jack

→ 2-loop finiteness

Jones

## Reduction of couplings

• Extension of method in SSB sector

+ application in min susy SU(5)

Kubo  
Mondragon  
Z

•• 1-loop sum rule for soft

Kawanuma

scalar masses in non-finite

Kobayashi

Kubo

susy ths.

••• 2-loop sum rule for soft

Kobayashi

Kubo

scalar masses in finite ths.

Mondragon  
Z

\* All-loop RGI relations

Yamada

Hisano,

Shifman

in finite and non-finite ths

Kazakov

Jack, Jones,

Pickering

\* \* All-loop sum rule for  
soft scalar masses in finite  
and non-finite t.h.s

Kobayashi  
Kubo  
2

• • SU(5) FUTs

Kobayashi  
Kubo  
Mondragon  
2

• Prediction of  $s$ -spectrum in  
terms of few parameters starting  
from several hundreds GeV.

• • The LSP is neutralino ✓ (see e.g.  
Kazakov  
et. al.  
Yoshioka)

• • Radiative E-W breaking ✓ (see e.g.  
Bignole  
Ibanez, Munoz)

• • • No funny colour, charge ✓ (see e.g.  
Casas et. al.)

\* Prediction of Higgs mass.

Lightest  $\sim 118 - 129$  GeV

Similar results also for min susy SU(5)

Consider a chiral, anomaly free,  
 $N=1$  gauge theory with group  $G$ .

The superpotential is

$$W = \frac{1}{6} Y^{ijk} \Phi_i \Phi_j \Phi_k + \frac{1}{2} \mu^{ij} \Phi_i \Phi_j$$

$\left. \begin{array}{l} Y^{ijk} \\ \mu^{ij} \end{array} \right\}$  gauge invariant  
Yukawa couplings

$\Phi_i$  - matter superfields  
in irreducible reps of  $G$

Necessary and sufficient conditions  
for  $N=1$  1-loop finiteness:

- Vanishing of  $\beta_g^{(1)}$  implies

$$\sum_i \ell(R_i) = 3 C_2(G) \quad ||$$

$\ell(R_i)$  - Dynkin index of  $R_i$

$C_2(G)$  - Quadratic Casimir of  $G$  (adjoint)

$\Rightarrow$  Selection of the field content  
(representations) of the theory

• • Vanishing of  $\gamma^i_j$  implies

$$\begin{array}{ccc} Y^{ikl} Y_{jkl} = 2 \delta^i_j g^2 C_2(R_i) // \\ \uparrow & & \uparrow \\ \text{Yukawa} & & \text{gauge} \end{array}$$

$C_2(R_i)$  - quadratic Casimir of  $R_i$

$$Y_{ijk} = (Y_{ijk})^*$$

⇒ Yukawa and gauge couplings are related.

Note •  $\mu^i$ s are not restricted

• • Appearance of  $U(1)$  is incompatible with 1<sup>st</sup> cond.

• • 2<sup>nd</sup> cond forbids the presence of singlets with nonvanishing couplings.

∴ ⇒ ~~Susy~~ by  $G$ -invariant soft terms

\* 1-loop finiteness condts necessary and sufficient to guarantee 2-loop finiteness

\* 1-loop finiteness condts ensure that  $\beta_g^{(3)}$  in 3-loops vanishes but in general  $\gamma^{(3)}$  does not.

Grisaru - Milewski - Zanon

Parke - West

What happens in higher loops?

So far 1-loop finiteness condts (on  $\gamma_s$ ) are telling us

$$\gamma^{ijk} = \gamma^{ijk}(g)$$

$$\beta_{\gamma}^{(1)ijk} = 0$$

\* \* Necessary and sufficient condt  
for vanishing  $b_g$  and  $b_{ijk}$  to all  
orders

1.  $b_g^{(1)} = 0$

Lucchesi  
Piquet  
Sibold

2.  $\gamma_s^{(1)i} = 0$

3.  $b_Y^{ijk} = b_g \frac{dY^{ijk}}{dg}$

admit power series solution which  
in lowest order is a solution of  
condt 2.

3'. There exist solutions to  $\gamma_s^{(1)i} = 0$

of the form

$Y^{ijk} = p^{ijk} g$ ,  $p^{ijk}$ -complex

4. These solutions are isolated  
and non-degenerate considered  
as solutions of  $b_Y^{(1)ijk} = 0$



Recall

R-invariance, axial anomaly

In massless  $N=1$  ths

$U(1)$  chiral transformation  $R$ :

$$A_\mu \rightarrow A_\mu, \quad f \rightarrow e^{-i\alpha} f,$$

$$\phi \rightarrow e^{-i\frac{2}{3}\alpha} \phi, \quad \psi \rightarrow e^{i\frac{1}{3}\alpha} \psi, \dots$$

$$\Psi_D = \begin{pmatrix} \psi \\ \bar{\chi} \end{pmatrix} \rightarrow e^{i\alpha\gamma_5} \Psi_D$$

Noether current  $J_R^\mu = \bar{\lambda}_D \gamma^\mu \gamma^5 \lambda_D + \dots$

$$\leadsto \partial_\mu J_R^\mu = r (\epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma} + \dots)$$

$$r = \frac{1}{16\pi^2} g^2$$

Only 1-loop contributions

due to non-renormalization thm.

Adler, Bardeen, Jackiw, Pi, Shei, Zee

# Supercurrent

$$\mathcal{J} \equiv \left\{ \underset{\substack{\text{associated} \\ \text{to } P\text{-invariance}}}{J_R^{\mu\nu}}, \underset{\substack{\text{associated} \\ \text{to SUSY}}}{Q_\alpha^\mu}, \underset{\substack{\text{associated} \\ \text{to translations}}}{T_\nu^\mu} \right\}, \dots \text{vector supermultiplet}$$

Ferrara + Zumino

(supercurrent is represented as vector superfield)

$$V_\mu(x, \theta, \bar{\theta}) = R_\mu(x) - i \theta^\alpha Q_{\mu\alpha}(x) + i \bar{\theta}_{\dot{\alpha}} \bar{Q}_{\mu}^{\dot{\alpha}}(x) - 2(\theta\sigma^\nu\bar{\theta}) T_{\mu\nu}(x) + \dots$$

- $J_R^{\mu\nu} \neq J_R^\mu$
- $J_R^{\mu\nu} = J_R^\mu + O(\theta)$

In addition

(Clarke, Piquet, Sibold)

$$\mathcal{J} = \left\{ b_g F^{\mu\nu} F_{\mu\nu} + \dots, b_g \in \text{MPG } F_{\mu\nu} F_{\rho\sigma}, \dots \right\}$$

Super-trace anomaly at  $T_\nu^\mu$  anomaly!  $\rightarrow$  R-current

$$\left. \left\{ b_g \int \sigma_{\alpha\beta}^{\mu\nu} F_{\mu\nu} + \dots, \dots \right\} \right\} \text{chiral supermultiplet}$$

trace anomaly of susy current

There is a relation, whose structure is independent from the renormalization scheme, although individual coefficients (except the 1-loop values of  $\beta$ -functions) may be scheme dependent

$$r = b_g (1 + x_g) + B_{ijk} x^{ijk} - \gamma_A r^A$$

radiative corrections

linear combinations of anomalous dims

unrenormalized coefficients of anomalies associated to chiral inv. of superpotential

Then: (i) no gauge anomaly

(ii)  $\beta''(g) = 0$  i.e. no R-current anomaly

(iii)  $\gamma^{(1)j} = 0$  implies also  $r^A = 0$

(iv) exist solutions to  $\gamma^{(1)} = 0$  of the

form  $C_{ijk} = P_{ijk} g$ ,  $P_{ijk}$  - complex

(v) these solutions are isolated + non-degenerate

when considered as solutions of  $B_{ijk}^{(1)} = 0$

- Then each of the solutions can be uniquely extended to a formal power series in  $g$ , and the  $N=1$  Y-M models depend on the single coupling constant  $g$  with a  $\beta$ -function vanishing to all orders.

Proof: Inserting  $B_{ijk} = b_g \frac{dB_{ijk}}{dg}$  in the identity and taking into account the vanishing of  $r, r^A$

$$\leadsto 0 = b_g (1 + O(\hbar))$$

Its solution (as formal power series in  $\hbar$ ) is:  $b_g = 0$   
and  $B_{ijk} = 0$  too.  $\parallel$

# 2-loop RGEs for SSB parameters

Martin-Vaughn - Yamada - Jack - Jones  
1994

Consider  $N=1$  gauge theory with

$$W = \frac{1}{6} Y^{ijk} \Phi_i \Phi_j \Phi_k + \frac{1}{2} \mu^{ij} \Phi_i \Phi_j$$

and SSB terms

$$-\mathcal{L}_{SSB} = \frac{1}{6} h^{ijk} \phi_i \phi_j \phi_k + \frac{1}{2} b^{ij} \phi_i \phi_j$$

$$+ \frac{1}{2} (m^2)_j^i \phi^{*i} \phi_j + \frac{1}{2} M \lambda \lambda + \text{h.c.}$$

• 1-loop finiteness conditions

$$h^{ijk} = -M Y^{ijk}$$

$$(m^2)_j^i = \frac{1}{3} M M^* \delta_j^i \quad \text{universality}$$

in addition to  $\beta_g^{(1)} = \gamma^{(1)i}_j = 0$

• • 1-loop finiteness

$\leadsto$  2-loop finiteness

Assuming

- $b_g^{(1)} = \gamma^{(1)j}_i = 0$

- the reduction eq

$$b_Y^{ijk} = b_g d \gamma^{ijk} / dg$$

admits power series solution

$$\gamma^{ijk} = g \sum_{n=0} P_{(n)}^{ijk} g^{2n}$$

- $(m^2)_j^i = m_j^2 \delta_j^i$

$$\Rightarrow (m_i^2 + m_j^2 + m_k^2) / MM^* = 1 + \frac{g^2}{16\pi^2} \Delta^{(2)}$$

for  $i, j, k$  with  $P_{(10)}^{ijk} \neq 0$

where  $\Delta^{(2)} = -2 \sum_l \left[ (m_l^2 / MM^*) - \frac{1}{3} \right] \ell(\ell)$

- $\Delta^{(2)} = 0$  for  $N=4$  with 5Tr cond

- $\Delta^{(2)} = 0$  for the  $N=1, SU(5)$  FUTs!

# The $SU(5)$ finite model

Kapetanakis, Mondragon, Z

Kobayashi, Kubo, Mondragon, Z

<u>Content</u>	$H_\alpha$	$\bar{H}_\alpha$	
$3(\bar{5} + 10) + 4(5 + \bar{5}) + 24$			Jones-Raby Hamidi-Schwartz Garcia et al Kazakov Babu-Enkhbaatar Gogoladze
↑ fermion supermultiplets		← scalar supermultiplets	

$$\Rightarrow W = \sum_{i=1}^3 \left[ \frac{1}{2} g_i^u 10_i 10_i H_i + g_i^d 10_i \bar{5}_i \bar{H}_i \right]$$

$$+ g_{23}^u 10_2 10_3 H_4 + g_{23}^d 10_2 \bar{5}_3 \bar{H}_4 + g_{32}^d 10_3 \bar{5}_2 \bar{H}_4$$

$$+ \sum_{\alpha=1}^4 g_\alpha^f H_\alpha 24 \bar{H}_\alpha + g' / 3 (24)^3$$

(with enhanced discrete symmetry  
after reduction of couplings)

We find

$$b_g^{(1)} = 0$$

$$b_{i\alpha}^{u(1)} = \frac{1}{16\pi^2} \left[ -\frac{96}{5} g^2 + \sum_{b=1}^4 (g_{ib}^u)^2 + 3 \sum_{j=1}^3 (g_{ja}^u)^2 + \frac{24}{5} (g_\alpha^f)^2 + 4 \sum_{b=1}^4 (g_{ib}^d)^2 \right] g_{i\alpha}^u$$

$$b_{i\alpha}^{d(1)} = \frac{1}{16\pi^2} \left[ -\frac{84}{5} g^2 + 3 \sum_{b=1}^4 (g_{ib}^u)^2 + \frac{24}{5} (g_\alpha^f)^2 + 4 \sum_{j=1}^3 (g_{ja}^d)^2 + 6 \sum_{b=1}^4 (g_{ib}^d)^2 \right] g_{i\alpha}^d$$

$$b_{i\alpha}^{\lambda(1)} = \frac{1}{16\pi^2} \left[ -30 g^2 + \frac{63}{5} (g^\lambda)^2 + 3 \sum_{\alpha=1}^4 (g_\alpha^f)^2 \right] g_{i\alpha}^\lambda$$

$$b_\alpha^{f(1)} = \frac{1}{16\pi^2} \left[ -\frac{98}{5} g^2 + 3 \sum_{i=1}^3 (g_{i\alpha}^u)^2 \right. \\ \left. + 4 \sum_{i=1}^3 (g_{i\alpha}^d)^2 + \frac{48}{5} (g_\alpha^f)^2 \right. \\ \left. + \sum_{b=1}^4 (g_b^f)^2 + \frac{21}{5} (g^\lambda)^2 \right] g_\alpha^f$$



Considering  $g$  as the primary coupling, we solve the Ren. Eqs.

$$\beta_g = \beta_a \frac{dg}{d\ln a}$$

requiring power series ansatz.

$$\Rightarrow (g_{ii}^u)^2 = \frac{8}{5} g^2 + \dots, (g_{ii}^d)^2 = \frac{6}{5} g^2 + \dots$$

$$(g^x)^2 = \frac{15}{7} g^2 + \dots, (g_4^f)^2 = g^2, (g_\alpha^f)^2 = 0 + \dots (\alpha=1,2,3)$$

Higher order terms can be ~~minimally~~  
by determined.

$\Rightarrow$  All 1-loop  $\beta$ -functions  
vanish.

Moreover

All 1-loop anomalous dimensions of chiral fields vanish.

$$\gamma_{10i}^{(1)} = \frac{1}{16\pi^2} \left[ -\frac{36}{5} g^2 + 3 \sum_{b=1}^4 (g_{ib}^u)^2 + 2 \sum_{b=1}^4 (g_{ib}^d)^2 \right]$$

$$\gamma_{\bar{5}i}^{(1)} = \frac{1}{16\pi^2} \left[ -\frac{24}{5} g^2 + 4 \sum_{b=1}^4 (g_{ib}^d)^2 \right]$$

$$\gamma_{H\alpha}^{(1)} = \frac{1}{16\pi^2} \left[ -\frac{24}{5} g^2 + 3 \sum_{i=1}^3 (g_{i\alpha}^u)^2 + \frac{24}{5} (g_\alpha^f)^2 \right]$$

$$\gamma_{\bar{H}\alpha}^{(1)} = \frac{1}{16\pi^2} \left[ -\frac{24}{5} g^2 + 4 \sum_{i=1}^3 (g_{i\alpha}^d)^2 + \frac{24}{5} (g_\alpha^f)^2 \right]$$

$$\gamma_{24}^{(1)} = \frac{1}{16\pi^2} \left[ -\frac{10}{5} g^2 + \sum_{\alpha=1}^4 (g_\alpha^f)^2 + \frac{21}{5} (g^\lambda)^2 \right]$$

$\Rightarrow$  Necessary and sufficient conditions for finiteness to all orders are satisfied.

- $SU(5)$  breaks down to the standard model due to  $\langle 24 \rangle$
- Use the freedom in mass parameters to obtain only a pair of Higgs fields light, acquiring v.e.v.
- Get rid of unwanted triplets rotating the Higgs sector (after a fine tuning)  
see Quiros et. al., Kazakov et. al  
Yoshioka
- Adding soft terms we can achieve SUSY breaking.

1) Gauge Couplings Unification  
 $\sin^2 \theta_w, \alpha_{em} \rightarrow \alpha_3(M_Z)$  Marciano+Serjenuic  
 Anibaldi et. al.

2) Bottom-Tau Yukawa Unif.  
 SU(5) type  
 $\rightarrow m_t \sim 100 - 200 \text{ GeV}$  Barger et. al.  
 Carena et. al.

\*3) Top-Bottom-Tau Yuk. Unif.  
 $h_t^2 = \frac{4}{3} h_{b,T}^2$  in SU(5) type  
 Similar to SU(5) Ananthanarayan et. al.  
 Barger et. al.  
 $\rightarrow m_t \sim 160 - 200 \text{ GeV}$  Carena et. al.

\*4) Gauge-Top-Bottom-Tau Unif.  
 e.g. SU(5):  $h_t^2 = \frac{8}{5} g_U^2$ ;  $h_{b,T}^2 = \frac{6}{5} g_U^2$

$M_s$ [GeV]	$\alpha_{3(5f)}(M_Z)$	$\tan \beta$	$M_{GUT}$ [GeV]	$M_b$ [GeV]	$M_t$ [GeV]
300	0.123	54.1	$2.2 \times 10^{16}$	5.3	183
500	0.122	54.2	$1.9 \times 10^{16}$	5.3	183
$10^3$	0.120	54.3	$1.5 \times 10^{16}$	5.2	184

**FUTA**

$M_s$ [GeV]	$\alpha_{3(5f)}(M_Z)$	$\tan \beta$	$M_{GUT}$ [GeV]	$M_b$ [GeV]	$M_t$ [GeV]
800	0.120	48.2	$1.5 \times 10^{16}$	5.4	174
$10^3$	0.119	48.2	$1.4 \times 10^{16}$	5.4	174
$1.2 \times 10^3$	0.118	48.2	$1.3 \times 10^{16}$	5.4	174

**FUTB**

$M_s$ [GeV]	$\alpha_{3(5f)}(M_Z)$	$\tan \beta$	$M_{GUT}$ [GeV]	$M_b$ [GeV]	$M_t$ [GeV]
300	0.123	47.9	$2.2 \times 10^{16}$	5.5	178
500	0.122	47.8	$1.8 \times 10^{16}$	5.4	178
1000	0.119	47.7	$1.5 \times 10^{16}$	5.4	178

**MIN SU(5)**

The predictions for the three models for different  $M_s$

*With theoretical corrections and uncertainties*


$\sim 4\%$

$M_t = 173.8 \pm 5 \text{ GeV}$      $178.0 \pm 4.3 \text{ GeV}$

CDF + D0

## Model A


Similar behaviour holds for Model B too



$m_{\tilde{\tau}}^2$  and  $m_\chi^2$  for the universal choice of soft scalar masses


# Model A

$M_{\text{susy}} = 0.3 \text{ TeV}$



The empty region yields a neutralino as LSP


Application of  $m_t^{\text{pole}}$ :



- ⇒ FUTB gives the correct prediction for  $m_t$
- ⇒ FUTA is ruled out experimentally




Application of  $m_b(M_Z)$ :



$\Rightarrow \mu < 0$  strongly favored

$\Rightarrow \mu > 0$  experimentally excluded

### 3D) Predictions for the light Higgs boson



green: consistent with  $B$  physics constraints

red: agreement with (loose) CDM bound

$118 \text{ GeV} \leq M_h \leq 129 \text{ GeV}$  (incl. theor. unc.)

$\Rightarrow$  "easy" to find for LHC (but "only" SM-like ...)

Typical mass spectrum for FUTB-:

$m_t$	172	$\overline{m_b}(M_Z)$	2.7
$\tan \beta =$	46	$\alpha_s$	0.116
$m_{\tilde{\chi}_1^0}$	796	$m_{\tilde{\tau}_2}$	1268
$m_{\tilde{\chi}_2^0}$	1462	$m_{\tilde{\nu}_3}$	1575
$m_{\tilde{\chi}_3^0}$	2048	$\mu$	-2046
$m_{\tilde{\chi}_4^0}$	2052	$B$	4722
$m_{\tilde{\chi}_1^\pm}$	1462	$M_A$	870
$m_{\tilde{\chi}_2^\pm}$	2052	$M_{H^\pm}$	875
$m_{\tilde{\tau}_1}$	2478	$M_H$	869
$m_{\tilde{\tau}_2}$	2804	$M_h$	124
$m_{\tilde{b}_1}$	2513	$M_1$	796
$m_{\tilde{b}_2}$	2783	$M_2$	1467
$m_{\tilde{\tau}_1}$	798	$M_3$	3655