

Some T-dual bundles over noncommutative spaces

Giovanni Landi

Trieste

Quantum spacetime structures: Dualities and new geometries

Bayrischzell, April 29- May 3, 2016

Work done over few years with

Francesca Arici

Simon Brain

Francesco D'Andrea

Jens Kaad

Abstract:

- Pimsner algebras of ‘tautological’ line bundles:

A procedure to construct total spaces of principal bundles out of a Fock-space construction

- Gysin-like sequences in KK-theory (D-brane charges)
- some hint to T-dual noncommutative bundles
- Examples: Quantum **lens spaces** as direct sums of line bundles over **weighted quantum projective spaces**

'grand motivations' :

Gauge fields on noncommutative spaces

T-duality for noncommutative spaces

Chern-Simons theory

A Gysin sequence for $U(1)$ -bundles

relates H -flux (three-forms on the total space E) to line bundles (two-forms on the base space M) also giving an isomorphism between Dixmier-Douady classes on E and line bundles on M

The classical Gysin sequence

Long exact sequence in cohomology; for any sphere bundle

In particular, for circle bundles: $U(1) \rightarrow E \xrightarrow{\pi} X$

$$\dots \longrightarrow H^k(E) \xrightarrow{\pi_*} H^{k-1}(X) \xrightarrow{\cup c_1(E)} H^{k+1}(X) \xrightarrow{\pi^*} H^{k+1}(E) \longrightarrow \dots$$

$$\dots \longrightarrow H^3(M) \xrightarrow{\pi^*} H^3(E) \xrightarrow{\pi_*} H^2(X) \xrightarrow{\cup c_1(E)} H^4(X) \longrightarrow \dots$$

$$H^3(E) \ni H \mapsto \pi_*(H) = F' = c_1(E')$$

$$\begin{array}{ccc}
 & E \times_M E' & \\
 & \swarrow \quad \searrow & \\
 E & & E' \\
 & \searrow \quad \swarrow & \\
 & M &
 \end{array}$$

π (arrow from E to M) π' (arrow from E' to M)

$$\dots \longrightarrow H^3(M) \xrightarrow{\pi^*} H^3(E') \xrightarrow{\pi_*} H^2(X) \xrightarrow{\cup c_1(E')} H^4(X) \longrightarrow \dots$$

$$F' \cup F = 0 = F \cup F'$$

$$\Rightarrow \exists H^3(E') \ni H' \mapsto \pi_*(H) = F = c_1(E)$$

T-dual (E, H) and (E', H')

Bouwknegt, Evslin, Mathai, 2004

difficult to generalize to quantum spaces

rather go to K-theory ; a six term exact sequence (see later)

K-theory elements = D-branes charges

Projective spaces and lens spaces

$$\mathbb{C}P^n = S^{2n+1}/U(1) \quad \text{and} \quad L^{(n,r)} = S^{2n+1}/\mathbb{Z}_r$$

assemble in principal bundles : $S^{2n+1} \longrightarrow L^{(n,r)} \xrightarrow{\pi} \mathbb{C}P^n$

This leads to the **Gysin sequence** in topological K-theory:

$$0 \longrightarrow K^1(L^{(n,r)}) \xrightarrow{\delta} K^0(\mathbb{C}P^n) \xrightarrow{\alpha} K^0(\mathbb{C}P^n) \xrightarrow{\pi^*} K^0(L^{(n,r)}) \longrightarrow 0$$

δ is a 'connecting homomorphism'

α is multiplication by the **Euler class** $\chi(\mathcal{L}_{-r}) := 1 - [\mathcal{L}_{-r}]$

From this:

$$K^1(L^{(n,r)}) \simeq \ker(\alpha) \quad \text{and} \quad K^0(L^{(n,r)}) \simeq \text{coker}(\alpha)$$

torsion groups

U(1)-principal bundles

The Hopf algebra

$$\mathcal{H} = \mathcal{O}(U(1)) := \mathbb{C}[z, z^{-1}] / \langle 1 - zz^{-1} \rangle$$

$$\Delta : z^n \mapsto z^n \otimes z^n \quad ; \quad S : z^n \mapsto z^{-n} \quad ; \quad \epsilon : z^n \mapsto 1$$

Let \mathcal{A} be a right comodule algebra over \mathcal{H} with coaction

$$\Delta_R : \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{H}$$

$\mathcal{B} := \{x \in \mathcal{A} \mid \Delta_R(x) = x \otimes 1\}$ be the subalgebra of coinvariants

Definition 1. *The datum $(\mathcal{A}, \mathcal{H}, \mathcal{B})$ is a quantum principal U(1)-bundle when the canonical map is an isomorphism*

$$\text{can} : \mathcal{A} \otimes_{\mathcal{B}} \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{H}, \quad x \otimes y \mapsto x \Delta_R(y).$$

\mathbb{Z} -graded algebras

$\mathcal{A} = \bigoplus_{n \in \mathbb{Z}} \mathcal{A}_n$ a \mathbb{Z} -graded algebra. A right \mathcal{H} -comodule algebra:

$$\Delta_R : \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{H} \quad x \mapsto x \otimes z^{-n}, \quad \text{for } x \in \mathcal{A}_n,$$

with the subalgebra of coinvariants given by \mathcal{A}_0 .

Proposition 2. *The triple $(\mathcal{A}, \mathcal{H}, \mathcal{A}_0)$ is a quantum principal $U(1)$ -bundle if and only if there exist finite sequences*

$$\{\xi_j\}_{j=1}^N, \{\beta_i\}_{i=1}^M \text{ in } \mathcal{A}_1 \quad \text{and} \quad \{\eta_j\}_{j=1}^N, \{\alpha_i\}_{i=1}^M \text{ in } \mathcal{A}_{-1}$$

such that:

$$\sum_{j=1}^N \xi_j \eta_j = 1_{\mathcal{A}} = \sum_{i=1}^M \alpha_i \beta_i.$$

Corollary 3. *Same conditions as above. The right-modules \mathcal{A}_1 and \mathcal{A}_{-1} are finitely generated and projective over \mathcal{A}_0 .*

Proof. For \mathcal{A}_1 : define the module homomorphisms

$$\Phi_1 : \mathcal{A}_1 \rightarrow (\mathcal{A}_0)^N, \quad \Phi_1(\zeta) = \begin{pmatrix} \eta_1 \zeta \\ \eta_2 \zeta \\ \vdots \\ \eta_N \zeta \end{pmatrix} \quad \text{and}$$
$$\Psi_1 : (\mathcal{A}_0)^N \rightarrow \mathcal{A}_1, \quad \Psi_1 \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_N \end{pmatrix} = \sum_j \xi_j x_j.$$

Then $\Psi_1 \Phi_1 = \text{Id}_{\mathcal{A}_1}$.

Thus $E_1 := \Phi_1 \Psi_1$ is an idempotent in $M_N(\mathcal{A}_0)$. □

The above results show that $(\mathcal{A}, \mathcal{H}, \mathcal{A}_0)$ is a quantum principal U(1)-bundle if and only if \mathcal{A} is *strongly \mathbb{Z} -graded*, that is

$$\mathcal{A}_n \mathcal{A}_m = \mathcal{A}_{n+m}$$

Equivalently, the right-modules $\mathcal{A}_{(\pm 1)}$ are finitely generated and projective over \mathcal{A}_0 if and only if \mathcal{A} is *strongly \mathbb{Z} -graded*

C. Nastasescu, F. Van Oystaeyen, *Graded Ring Theory*

K.H. Ulbrich, 1981

More generally: G any group with unit e

An algebra \mathcal{A} is G -graded if $\mathcal{A} = \bigoplus_{g \in G} \mathcal{A}_g$, and $\mathcal{A}_g \mathcal{A}_h \subseteq \mathcal{A}_{gh}$

If $\mathcal{H} := \mathbb{C}G$ the group algebra, then \mathcal{A} is G -graded if and only if \mathcal{A} is a right \mathcal{H} -comodule algebra for the coaction $\delta : \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{H}$

$$\delta(a_g) = a_g \otimes g, \quad a_g \in \mathcal{A}_g;$$

coinvariants given by $\mathcal{A}^{co\mathcal{H}} = \mathcal{A}_e$, the identity components.

Proposition 4. *The datum $(\mathcal{A}, \mathcal{H}, \mathcal{A}_e)$ is a noncommutative principal \mathcal{H} -bundle for the canonical map*

$$\text{can} : \mathcal{A} \otimes_{\mathcal{A}_e} \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{H}, \quad a \otimes b \mapsto \sum_g ab_g \otimes g,$$

if and only if \mathcal{A} is strongly graded, that is $\mathcal{A}_g \mathcal{A}_h = \mathcal{A}_{gh}$.

When $G = \mathbb{Z} = \widehat{U(1)}$, then $\mathbb{C}G = \mathcal{O}(U(1))$ as before.

More general scheme: Pimsner algebras M.V. Pimsner '97

The right-modules \mathcal{A}_1 and \mathcal{A}_{-1} before are 'line bundles' over \mathcal{A}_0

The slogan:

a line bundle

is a

self-Morita equivalence bimodule

Morita equivalence in one-page

E a (right) Hilbert module over B

B -valued hermitian structure $\langle \cdot, \cdot \rangle_\bullet$ on E

$\mathcal{L}(E)$ adjointable operators; $\mathcal{K}(E) \subseteq \mathcal{L}(E)$ compact operators

With $\xi, \eta \in E$, denote $\theta_{\xi, \eta} \in \mathcal{K}(E)$ defined by $\theta_{\xi, \eta}(\zeta) = \xi \langle \eta, \zeta \rangle$

$\mathcal{K}(E)$ -valued hermitian structure $\bullet \langle \cdot, \cdot \rangle$ on E : $\bullet \langle \xi, \eta \rangle := \theta_{\xi, \eta}$

The hermitian structures are compatible by construction

$$\bullet \langle \xi, \eta \rangle \zeta = \xi \langle \eta, \zeta \rangle_\bullet$$

Algebras $\mathcal{K}(E)$ and B are **Morita equivalent** via the bimodule E .

For a line bundle we are asking that there is an isomorphism $\phi : B \rightarrow \mathcal{K}(E)$ and thus E is a **B -bimodule**

Comparing with before:

$$\mathcal{A}_0 \rightsquigarrow B \quad \text{and} \quad \mathcal{A}_{-1} \rightsquigarrow E$$

Look for the analogue of $\mathcal{A} \rightsquigarrow \mathcal{O}_E$ Pimsner algebra

Examples

$$B = \mathcal{O}(\mathbb{C}P_q^n) \quad \text{quantum (weighted) projective spaces}$$

$$E = \mathcal{L}_{-r} \simeq (\mathcal{L}_{-1})^r \quad \text{(powers of) tautological line bundle}$$

$$\mathcal{O}_E = \mathcal{O}(\mathcal{L}_q^{(n,r)}) \quad \text{quantum lens spaces}$$

Define the B -module

$$E_\infty := \bigoplus_{N \in \mathbb{Z}} E^{\widehat{\otimes}_\phi N}, \quad E^0 = B$$

$E \otimes_\phi E$ the inner tensor product: a B -Hilbert module with B -valued hermitian structure

$$\langle \xi_1 \otimes \eta_1, \xi_2 \otimes \eta_2 \rangle = \langle \eta_1, \phi(\langle \xi_1, \xi_2 \rangle) \eta_2 \rangle$$

$E^{-1} = E^*$ the dual module;

its elements are written as λ_ξ for $\xi \in E$: $\lambda_\xi(\eta) = \langle \xi, \eta \rangle$

For each $\xi \in E$ a bounded adjointable operator

$$S_\xi : E_\infty \rightarrow E_\infty$$

generated by $S_\xi : E^{\widehat{\otimes}_\phi N} \rightarrow E^{\widehat{\otimes}_\phi (N+1)}$:

$$\begin{aligned} S_\xi(b) &:= \xi b, & b \in B, \\ S_\xi(\xi_1 \otimes \cdots \otimes \xi_N) &:= \xi \otimes \xi_1 \otimes \cdots \otimes \xi_N, & N > 0, \\ S_\xi(\lambda_{\xi_1} \otimes \cdots \otimes \lambda_{\xi_{-N}}) &:= \lambda_{\xi_2 \phi^{-1}(\theta_{\xi_1, \xi})} \otimes \lambda_{\xi_3} \otimes \cdots \otimes \lambda_{\xi_{-N}}, & N < 0. \end{aligned}$$

Definition 5. The *Pimsner algebra* \mathcal{O}_E of the pair (ϕ, E) is the smallest subalgebra of $\mathcal{L}(E_\infty)$ which contains the operators $S_\xi : E_\infty \rightarrow E_\infty$ for all $\xi \in E$.

Pimsner: universality of \mathcal{O}_E

There is a natural inclusion

$$B \hookrightarrow \mathcal{O}_E \quad \text{a generalized principal circle bundle}$$

roughly: as a vector space $\mathcal{O}_E \simeq E_\infty$ and

$$E^{\widehat{\otimes} \phi^N} \ni \eta \mapsto \eta \lambda^{-N}, \quad \lambda \in \mathbf{U}(1)$$

Two natural classes in KK-theory:

1. the class $[E] \in KK_0(B, B)$
of the even Kasparov module $(E, \phi, 0)$ (with trivial grading)

the map $\mathbf{1} - [E]$ has the role of the *Euler class* $\chi(E) := \mathbf{1} - [E]$

of the line bundle E over the ‘noncommutative space’ B

2. the class $[\partial] \in KK_1(\mathcal{O}_E, B)$

of the odd Kasparov module $(E_\infty, \tilde{\phi}, F)$:

$F := 2P - 1 \in \mathcal{L}(E_\infty)$ of the projection $P : E_\infty \rightarrow E_\infty$ with

$$\text{Im}(P) = \left(\bigoplus_{N=0}^{\infty} E^{\hat{\otimes}_{\phi} N} \right) \subseteq E_\infty$$

and inclusion $\tilde{\phi} : \mathcal{O}_E \rightarrow \mathcal{L}(E_\infty)$.

The Kasparov product induces group homomorphisms

$$[E] : K_*(B) \rightarrow K_*(B), \quad [E] : K^*(B) \rightarrow K^*(B)$$

and

$$[\partial] : K_*(\mathcal{O}_E) \rightarrow K_{*+1}(B), \quad [\partial] : K^*(B) \rightarrow K^{*+1}(\mathcal{O}_E),$$

Associated six-terms exact sequences **Gysin sequences**:
in K-theory:

$$\begin{array}{ccccc}
 K_0(B) & \xrightarrow{1-[E]} & K_0(B) & \xrightarrow{i_*} & K_0(\mathcal{O}_E) \\
 \uparrow [\partial] & & & & \downarrow [\partial] \\
 K_1(\mathcal{O}_E) & \xleftarrow{i_*} & K_1(B) & \xleftarrow{1-[E]} & K_1(B)
 \end{array} ;$$

the corresponding one in K-homology:

$$\begin{array}{ccccc}
 K^0(B) & \xleftarrow{1-[E]} & K^0(B) & \xleftarrow{i^*} & K^0(\mathcal{O}_E) \\
 \downarrow [\partial] & & & & \uparrow [\partial] \\
 K^1(\mathcal{O}_E) & \xrightarrow{i^*} & K^1(B) & \xrightarrow{1-[E]} & K^1(B)
 \end{array} .$$

In fact in KK-theory

The quantum spheres and the projective spaces

The coordinate algebra $\mathcal{O}(S_q^{2n+1})$ of quantum **sphere** S_q^{2n+1} :
-algebra generated by $2n + 2$ elements $\{z_i, z_i^\}_{i=0, \dots, n}$ s.t.:

$$\begin{aligned} z_i z_j &= q^{-1} z_j z_i & 0 \leq i < j \leq n, \\ z_i^* z_j &= q z_j z_i^* & i \neq j, \\ [z_n^*, z_n] &= 0, \quad [z_i^*, z_i] = (1 - q^2) \sum_{j=i+1}^n z_j z_j^* & i = 0, \dots, n-1, \end{aligned}$$

and a sphere relation:

$$1 = z_0 z_0^* + z_1 z_1^* + \dots + z_n z_n^* .$$

L. Vaksman, Ya. Soibelman, 1991 ; M. Welk, 2000

The $*$ -subalgebra of $\mathcal{O}(S_q^{2n+1})$ generated by

$$p_{ij} := z_i^* z_j$$

coordinate algebra $\mathcal{O}(\mathbb{C}P_q^n)$ of the quantum **projective space** $\mathbb{C}P_q^n$

Invariant elements for the $U(1)$ -action on the algebra $\mathcal{O}(S_q^{2n+1})$:

$$(z_0, z_1, \dots, z_n) \mapsto (\lambda z_0, \lambda z_1, \dots, \lambda z_n), \quad \lambda \in U(1).$$

the fibration $S_q^{2n+1} \rightarrow \mathbb{C}P_q^n$ is a quantum $U(1)$ -principal bundle:

$$\mathcal{O}(\mathbb{C}P_q^n) = \mathcal{O}(S_q^{2n+1})^{U(1)} \hookrightarrow \mathcal{O}(S_q^{2n+1}).$$

The C^* -algebras $C(S_q^{2n+1})$ and $C(\mathbb{C}P_q^n)$ of continuous functions: completions of $\mathcal{O}(S_q^{2n+1})$ and $\mathcal{O}(\mathbb{C}P_q^n)$ in the universal C^* -norms

these are **graph algebras** [J.H. Hong, W. Szymański 2002](#)

$$\Rightarrow K_0(\mathbb{C}P_q^n) \simeq \mathbb{Z}^{n+1} \simeq K^0(C(\mathbb{C}P_q^n))$$

[F. D'Andrea, G. L. 2010](#)

Generators of the homology group $K^0(C(\mathbb{C}P_q^n))$ given explicitly as (classes of) even Fredholm modules

$$\mu_k = (\mathcal{O}(\mathbb{C}P_q^n), \mathcal{H}_{(k)}, \pi^{(k)}, \gamma_{(k)}, F_{(k)}), \quad \text{for } 0 \leq k \leq n.$$

Generators of the K-theory $K_0(\mathbb{C}P_q^n)$ also given explicitly as projections whose entries are polynomial functions:

line bundles & projections

For $N \in \mathbb{Z}$, vector-valued functions

$$\Psi_N := (\psi_{j_0, \dots, j_n}^N) \quad \text{s.t.} \quad \Psi_N^* \Psi_N = 1$$

$\Rightarrow P_N := \Psi_N \Psi_N^*$ is a **projection**:

$$P_N \in M_{d_N}(\mathcal{O}(\mathbb{C}P_q^n)), \quad d_N := \binom{|N| + n}{n},$$

Entries of P_N are $U(1)$ -invariant and so elements of $\mathcal{O}(\mathbb{C}P_q^n)$

Proposition 6. For all $N \in \mathbb{N}$ and for all $0 \leq k \leq n$ it holds that

$$\langle [\mu_k], [P_{-N}] \rangle := \text{Tr}_{\mathcal{H}_k}(\gamma_{(k)}(\pi^{(k)}(\text{Tr } P_{-N})) = \binom{N}{k},$$

$[\mu_0], \dots, [\mu_n]$ are generators of $K^0(C(\mathbb{C}P_q^n))$,

and $[P_0], \dots, [P_{-n}]$ are generators of $K_0(\mathbb{C}P_q^n)$

The matrix of couplings $M \in M_{n+1}(\mathbb{Z})$ is invertible over \mathbb{Z} :

$$M_{ij} := \langle [\mu_i], [P_{-j}] \rangle = \binom{j}{i}, \quad (M^{-1})_{ij} = (-1)^{i+j} \binom{j}{i}.$$

These are bases of \mathbb{Z}^{n+1} as \mathbb{Z} -modules;

they generate \mathbb{Z}^{n+1} as an Abelian group.

The inclusion $\mathcal{O}(\mathbb{C}\mathbb{P}_q^n) \hookrightarrow \mathcal{O}(S_q^{2n+1})$ is a $U(1)$ q.p.b.

To a projection P_N there corresponds an **associated line bundle**

$$\mathcal{L}_N \simeq (\mathcal{O}(\mathbb{C}\mathbb{P}_q^n))^{d_N} P_N \simeq P_{-N} (\mathcal{O}(\mathbb{C}\mathbb{P}_q^n))^{d_N}$$

\mathcal{L}_N made of elements of $\mathcal{O}(S_q^{2n+1})$ transforming under $U(1)$ as

$$\varphi_N \mapsto \varphi_N \lambda^{-N}, \quad \lambda \in U(1)$$

Each \mathcal{L}_N is indeed a bimodule over $\mathcal{L}_0 = \mathcal{O}(\mathbb{C}\mathbb{P}_q^n)$; – **the bimodule of equivariant maps** for the IRREP of $U(1)$ with **weight N** . Also,

$$\mathcal{L}_N \otimes_{\mathcal{O}(\mathbb{C}\mathbb{P}_q^n)} \mathcal{L}_M \simeq \mathcal{L}_{N+M}$$

The module \mathcal{L}_N is a **line bundle**, in the sense that its '**rank**' (as computed by pairing with $[\mu_0]$) is equal to 1

Completely characterized by its '**first Chern number**' (as computed by pairing with the class $[\mu_1]$):

Proposition 7. *For all $N \in \mathbb{Z}$ it holds that*

$$\langle [\mu_0], [\mathcal{L}_N] \rangle = 1 \quad \text{and} \quad \langle [\mu_1], [\mathcal{L}_N] \rangle = -N.$$

The line bundle \mathcal{L}_{-1} emerges as a central character:
its only non-vanishing charges are

$$\langle [\mu_0], [\mathcal{L}_{-1}] \rangle = 1 \qquad \langle [\mu_1], [\mathcal{L}_{-1}] \rangle = 1$$

\mathcal{L}_{-1} is the *tautological line bundle* for $\mathbb{C}P_q^n$,

with *Euler class*

$$u = \chi([\mathcal{L}_{-1}]) := 1 - [\mathcal{L}_{-1}].$$

Proposition 8. *It holds that*

$$K_0(\mathbb{C}P_q^n) \simeq \mathbb{Z}[u]/u^{n+1} \simeq \mathbb{Z}^{n+1}.$$

$[\mu_k]$ and $(-u)^j$ are *dual bases* of K-homology and K-theory

The quantum lens spaces

Fix an integer $r \geq 2$ and define

$$\mathcal{O}(\mathbb{L}_q^{(n,r)}) := \bigoplus_{N \in \mathbb{Z}} \mathcal{L}_{rN}.$$

Proposition 9.

$\mathcal{O}(\mathbb{L}_q^{(n,r)})$ is a $*$ -algebra; all elements of $\mathcal{O}(S_q^{2n+1})$ invariant under the action $\alpha_r : \mathbb{Z}_r \rightarrow \text{Aut}(\mathcal{O}(S_q^{2n+1}))$ of the cyclic group \mathbb{Z}_r :

$$(z_0, z_1, \dots, z_n) \mapsto (e^{2\pi i/r} z_0, e^{2\pi i/r} z_1, \dots, e^{2\pi i/r} z_n).$$

The 'dual' $\mathbb{L}_q^{(n,r)}$:

the *quantum lens space* of dimension $2n + 1$ (and index r)

There are algebra inclusions

$$j : \mathcal{O}(\mathbb{C}P_q^n) \hookrightarrow \mathcal{O}(\mathbb{L}_q^{(n,r)}) \hookrightarrow \mathcal{O}(S_q^{2n+1}).$$

Pulling back line bundles

Proposition 10. *The algebra inclusion $j : \mathcal{O}(\mathbb{C}P_q^n) \hookrightarrow \mathcal{O}(L_q^{(n,r)})$ is a quantum principal bundle with structure group $\tilde{U}(1) := U(1)/\mathbb{Z}_r$:*

$$\mathcal{O}(\mathbb{C}P_q^n) = \mathcal{O}(L_q^{(n,r)})^{\tilde{U}(1)}.$$

Then one can ‘pull-back’ line bundles from $\mathbb{C}P_q^n$ to $L_q^{(n,r)}$.

$$\begin{array}{ccc} \tilde{\mathcal{L}}_N & \xleftarrow{j^*} & \mathcal{L}_N \\ \downarrow \text{dotted} & & \downarrow \text{dotted} \\ \mathcal{O}(L_q^{(n,r)}) & \xleftarrow{j} & \mathcal{O}(\mathbb{C}P_q^n). \end{array}$$

Definition 11. For each \mathcal{L}_N an $\mathcal{O}(\mathbb{C}P_q^n)$ -bimodule (a line bundle over $\mathbb{C}P_q^n$), its ‘pull-back’ to $L_q^{(n,r)}$ is the $\mathcal{O}(L_q^{(n,r)})$ -bimodule

$$\tilde{\mathcal{L}}_N = j_*(\mathcal{L}_N) := \mathcal{O}(L_q^{(n,r)}) \otimes_{\mathcal{O}(\mathbb{C}P_q^n)} \mathcal{L}_N.$$

The algebra inclusion $j : \mathcal{O}(\mathbb{C}P_q^n) \rightarrow \mathcal{O}(L_q^{(n,r)})$ induces a map

$$j_* : K_0(\mathbb{C}P_q^n) \rightarrow K_0(L_q^{(n,r)})$$

Each \mathcal{L}_N over $\mathbb{C}P_q^n$ is not free when $N \neq 0$,

this need not be the case for $\tilde{\mathcal{L}}_N$ over $L_q^{(n,r)}$:

the pull-back $\tilde{\mathcal{L}}_{-r}$ of \mathcal{L}_{-r} is tautologically free :

$$\tilde{\mathcal{L}}_{-r} = \mathcal{O}(L_q^{(n,r)}) \otimes_{\mathcal{L}_0} \mathcal{L}_{-r} \simeq \mathcal{O}(L_q^{(n,r)}) = \tilde{\mathcal{L}}_0.$$

$\Rightarrow (\tilde{\mathcal{L}}_{-N})^{\otimes r} \simeq \tilde{\mathcal{L}}_{-rN}$ also has trivial class for any $N \in \mathbb{Z}$

$\tilde{\mathcal{L}}_{-N}$ define *torsion classes*; they generate the group $K_0(L_q^{(n,r)})$

Multiplying by the Euler class

A second crucial ingredient

$$\alpha : K_0(\mathbb{C}P_q^n) \rightarrow K_0(\mathbb{C}P_q^n),$$

α is multiplication by $\chi(\mathcal{L}_{-r}) := 1 - [\mathcal{L}_{-r}]$

the **Euler class** of the line bundle \mathcal{L}_{-r}

Assembly these into an exact sequence, the *Gysin sequence*

$$0 \rightarrow K_1(\mathbb{L}_q^{(n,r)}) \xrightarrow{\partial} K_0(\mathbb{C}P_q^n) \xrightarrow{\alpha} K_0(\mathbb{C}P_q^n) \rightarrow K_0(\mathbb{L}_q^{(n,r)}) \rightarrow 0$$

$$0 \rightarrow K_1(\mathbb{L}_q^{(n,r)}) \xrightarrow{\text{Ind}_{\mathfrak{D}}} K_0(\mathbb{C}P_q^n) \rightarrow \dots$$

and

$$\dots \rightarrow K_0(\mathbb{L}_q^{(n,r)}) \xrightarrow{\text{Ind}_{\mathfrak{D}}} 0$$

$\text{Ind}_{\mathfrak{D}}$ comes from Kasparov theory

Some practical and important applications, notably, the computation of the K-theory of the quantum lens spaces $L_q^{(n,r)}$.

Thus

$$K_1(L_q^{(n,r)}) \simeq \ker(\alpha), \quad K_0(L_q^{(n,r)}) \simeq \operatorname{coker}(\alpha).$$

Moreover, *geometric* generators of the groups

$$K_1(L_q^{(n,r)}) \quad K_0(L_q^{(n,r)})$$

for the latter as pulled-back line bundles from $\mathbb{C}P_q^n$ to $L_q^{(n,r)}$

Explicit generators as integral combinations of powers of the pull-back to the lens space $L_q^{(n,r)}$ of the generator

$$u := 1 - [\mathcal{L}_{-1}]$$

The K-theory of quantum lens spaces

Proposition 12. *The $(n + 1) \times (n + 1)$ matrix α has rank n :*

$$K_1(C(L_q^{(n,r)})) \simeq \mathbb{Z}.$$

On the other hand, the structure of the **cokernel** of the matrix A depends on the divisibility properties of the integer r .

This leads to

$$K_0(L_q^{(n,r)}) = \mathbb{Z} \oplus \mathbb{Z}/\alpha_1\mathbb{Z} \oplus \cdots \oplus \mathbb{Z}/\alpha_n\mathbb{Z}.$$

for suitable integers $\alpha_1, \dots, \alpha_n$.

Example 13. For $n = 1$

$$K_0(C(L_q^{(1,r)})) = \mathbb{Z} \oplus \mathbb{Z}_r.$$

From definition $[\tilde{\mathcal{L}}_{-r}] = 1$, thus $\tilde{\mathcal{L}}_{-1}$ generates the torsion part.

Alternatively, from $u^2 = 0$ it follows that $\mathcal{L}_{-j} = -(j-1) + j\mathcal{L}_{-1}$; upon lifting to $L_q^{(1,r)}$, for $j = r$ this yields

$$r(1 - [\tilde{\mathcal{L}}_{-1}]) = 0$$

or $1 - [\tilde{\mathcal{L}}_{-1}]$ is cyclic of order r .

Example 14. If $r = 2$ $L_q^{(n,2)} = S_q^{2n+1}/\mathbb{Z}_2 = \mathbb{R}P_q^{2n+1}$,
the quantum real projective space, we get

$$K_0(C(\mathbb{R}P_q^{2n+1})) = \mathbb{Z} \oplus \mathbb{Z}_{2^n}$$

the generator $1 - [\tilde{\mathcal{L}}_{-1}]$ is cyclic with the correct order 2^n .

Example 15. For $n = 2$ there are two cases.

When $r = 2k + 1$:

$$r \tilde{u} = 0, \quad r \tilde{u}^2 = 0, \quad K_0(L_q^{(2,r)}) = \mathbb{Z} \oplus \mathbb{Z}_r \oplus \mathbb{Z}_r$$

When $r = 2k$:

$$\frac{1}{2}r (\tilde{u}^2 + 2\tilde{u}) = 0, \quad 2r \tilde{u} = 0, \quad K_0(C(L_q^{(2,r)})) = \mathbb{Z} \oplus \mathbb{Z}_{\frac{r}{2}} \oplus \mathbb{Z}_{2r}$$

T-dual Pimsner algebras: a simple example

$$0 \rightarrow K_1(\mathcal{L}_q^{(1,r)}) \xrightarrow{\partial} K_0(\mathbb{C}P_q^1) \xrightarrow{1 - [\mathcal{L}_{-r}]} K_0(\mathbb{C}P_q^1) \rightarrow K_0(\mathcal{L}_q^{(1,r)}) \rightarrow 0$$

$$\ker(1 - [\mathcal{L}_{-r}]) = \langle u \rangle = \langle 1 - [\mathcal{L}_{-1}] \rangle$$

\Rightarrow

$$K_1(\mathcal{L}_q^{(1,r)}) \ni h \mapsto \partial(h) = h(1 - [\mathcal{L}_{-1}]) \simeq 1 - [\mathcal{L}_{-h}]$$

and

$$(1 - [\mathcal{L}_{-r}])(1 - [\mathcal{L}_{-h}]) = 0 = (1 - [\mathcal{L}_{-h}])(1 - [\mathcal{L}_{-r}])$$

The exactness of the dual sequence for

$$0 \rightarrow K_1(\mathbb{L}_q^{(1,h)}) \xrightarrow{\partial} K_0(\mathbb{C}P_q^1) \xrightarrow{1 - [\mathcal{L}_{-h}]} K_0(\mathbb{C}P_q^1) \rightarrow K_0(\mathbb{L}_q^{(1,h)}) \rightarrow 0$$

implies there exists a $r \in K_1(\mathbb{L}_q^{(1,r)})$ such that

$$K_1(\mathbb{L}_q^{(1,h)}) \ni r \mapsto \partial(r) = r(1 - [\mathcal{L}_{-1}]) \simeq 1 - [\mathcal{L}_{-r}]$$

The couples

$$\left(\mathbb{L}_q^{(1,r)}, h \in K_1(\mathbb{L}_q^{(1,r)})\right) \text{ and } \left(\mathbb{L}_q^{(1,h)}, r \in K_1(\mathbb{L}_q^{(1,h)})\right)$$

are 'T-dual'

More generally : **Quantum w. projective lines and lens spaces:**

$B = \mathcal{O}(W_q(k, l)) =$ **quantum weighted projective line**

the fixed point algebra for a weighted circle action on $\mathcal{O}(S_q^3)$

$$z_0 \mapsto \lambda^k z_0, \quad z_1 \mapsto \lambda^l z_1, \quad \lambda \in U(1)$$

The corresponding universal enveloping C^* -algebra $C(W_q(k, l))$ does not in fact depend on the label k : isomorphic to the unitalization of l copies of $\mathcal{K} =$ compact operators on $l^2(\mathbb{N}_0)$

$$C(W_q(k, l)) = \widetilde{\bigoplus_{s=0}^l \mathcal{K}}$$

Then: $K_0(C(W_q(k, l))) = \mathbb{Z}^{l+1}, \quad K_1(C(W_q(k, l))) = 0$

a **partial resolution of singularities**, since classically

$$K_0(C(W(k, l))) = \mathbb{Z}^2.$$

$\mathcal{O}_E = \mathcal{O}(L_q(lk; k, l)) = \text{quantum lens space}$

Indeed, a vector space decomposition

$$\mathcal{O}(L_q(lk; k, l)) = \bigoplus_{N \in \mathbb{Z}} \mathcal{L}_N(k, l),$$

with $E = \mathcal{L}_1(k, l)$ a right finitely projective module

$$\mathcal{L}_1(k, l) := (z_1^*)^k \cdot \mathcal{O}(W_q(k, l)) + (z_0^*)^l \cdot \mathcal{O}(W_q(k, l))$$

Also, $\mathcal{O}(L_q(lk; k, l))$ the fixed point algebra of a cyclic action

$$\mathbb{Z}/(lk)\mathbb{Z} \times S_q^3 \rightarrow S_q^3$$

$$z_0 \mapsto \exp\left(\frac{2\pi i}{l}\right) z_0, \quad z_1 \mapsto \exp\left(\frac{2\pi i}{k}\right) z_1.$$

K-theory and K-homology of quantum lens space

Denote the diagonal inclusion by $\iota : \mathbb{Z} \rightarrow \mathbb{Z}^l$, $1 \mapsto (1, \dots, 1)$ with transpose $\iota^t : \mathbb{Z}^l \rightarrow \mathbb{Z}$, $\iota^t(m_1, \dots, m_l) = m_1 + \dots + m_l$.

Proposition 16. (Arici, Kaad, L.) With $k, l \in \mathbb{N}$ coprime:

$$\begin{aligned} K_0(L_q(lk; k, l)) &\simeq \text{coker}(1 - E) \simeq \mathbb{Z} \oplus (\mathbb{Z}^l / \text{Im}(\iota)) \\ K_1(L_q(lk; k, l)) &\simeq \ker(1 - E) \simeq \mathbb{Z}^l \end{aligned}$$

as well as

$$\begin{aligned} K^0(L_q(lk; k, l)) &\simeq \ker(1 - E^t) \simeq \mathbb{Z} \oplus (\ker(\iota^t)) \\ K^1(L_q(lk; k, l)) &\simeq \text{coker}(1 - E^t) \simeq \mathbb{Z}^l. \end{aligned}$$

Again there is no dependence on the label k .

Summing up:

A general procedure to construct principal circle bundle over a noncommutative space out of a Pimsner algebra connection

A Gysin like sequence relating D-brane charges.

consequences for

T-duality for noncommutative spaces

and

Chern-Simons theory

Thank you !!