

# Cosmology and particle physics

## Lecture notes

Timm Wrase

### Lecture 6 The thermal universe - part II

Last time we have seen that the standard model particles in the early universe were interacting so much that the Hubble expansion is negligible compared to the interaction rate, while the temperature is in the range  $100\text{GeV} \ll T \ll 10^{16}\text{GeV}$ . This means that there is ample time for the standard model particles to be in thermal equilibrium by the time the temperature is a few hundred  $\text{GeV}$ . We can therefore use equilibrium thermodynamics to discuss the evolution of this soup of standard model particles as the universe expands and cools.

#### 1 Thermal equilibrium

We also recalled last time that the phase space distribution function for particles in equilibrium (and with vanishing chemical potential) is given by

$$f_{\pm}(p) = \frac{1}{e^{E(p)/T} \pm 1}, \quad (1)$$

where the subscript  $+$  corresponds to fermions and the subscript  $-$  to bosons. In terms of this phase space distribution the number density  $n$ , the energy density  $\rho$  and the pressure  $P$  are given by

$$n \equiv \frac{g}{(2\pi)^3} \int d^3p f(p), \quad (2)$$

$$\rho \equiv \frac{g}{(2\pi)^3} \int d^3p f(p) E(p), \quad (3)$$

$$P \equiv \frac{g}{(2\pi)^3} \int d^3p f(p) \frac{p^2}{3E(p)}. \quad (4)$$

The integrals in  $n$ ,  $\rho$  and  $P$  have to be evaluated numerical unless we are in particular limits. For such limits we will make use of the general formulas

$$\int_0^{\infty} du \frac{u^n}{e^u - 1} = \zeta(n+1)\Gamma(n+1), \quad (5)$$

$$\int_0^{\infty} du u^n e^{-u^2} = \frac{1}{2}\Gamma\left(\frac{1}{2}(n+1)\right), \quad (6)$$

where  $\zeta$  is the Riemann zeta-function and the  $\Gamma$ -function is an extension of the factorial function and in particular takes the values  $\Gamma(n) = (n-1)!$  for  $n \in \mathbb{N}^*$ .

## 1.1 The relativistic limit

Let us first evaluate  $n$ ,  $\rho$  and  $P$  for relativistic particles:

$$E(p) = \sqrt{m^2 + p^2} \approx p \gg m. \quad (7)$$

We define  $y = p/T$  so that  $f_{\pm}(y) = 1/(e^y \pm 1)$ . For bosons we then find

$$n_b = \frac{g}{(2\pi)^3} \int_0^{\infty} \frac{4\pi T^3 y^2 dy}{e^y - 1} = \frac{gT^3 \zeta(3) \Gamma(3)}{2\pi^2} = \frac{\zeta(3)}{\pi^2} g T^3, \quad (8)$$

where  $\zeta(3) \approx 1.2$ . For fermions we can use that

$$\frac{1}{e^y + 1} = \frac{1}{e^y - 1} - \frac{2}{e^{2y} - 1}, \quad (9)$$

to get

$$n_f = n_b - \frac{g}{(2\pi)^3} \int_0^{\infty} \frac{8\pi T^3 y^2 dy}{e^{2y} - 1} = n_b - \frac{g}{(2\pi)^3} \int_0^{\infty} \frac{\pi T^3 \tilde{y}^2 d\tilde{y}}{e^{\tilde{y}} - 1} = n_b - \frac{1}{4} n_b = \frac{3\zeta(3)}{4\pi^2} g T^3. \quad (10)$$

So we have found for relativistic particles that

$$n_b = \frac{4}{3} n_f = \frac{\zeta(3)}{\pi^2} g T^3. \quad (11)$$

Now let us likewise calculate the energy density

$$\rho_b = \frac{g}{(2\pi)^3} \int_0^{\infty} \frac{4\pi T^4 y^3 dy}{e^y - 1} = \frac{g}{2\pi^2} T^4 \zeta(4) \Gamma(4) = \frac{\pi^2}{30} g T^4, \quad (12)$$

where we used that  $\zeta(4) = \pi^4/90$ . For fermions we find

$$\rho_f = \rho_b - \frac{g}{(2\pi)^3} \int_0^{\infty} \frac{8\pi T^4 y^3 dy}{e^{2y} - 1} = \rho_b - \frac{g}{(2\pi)^3} \int_0^{\infty} \frac{\pi T^4 \tilde{y}^3 d\tilde{y}}{e^{\tilde{y}} - 1} = \rho_b - \frac{1}{8} \rho_b = \frac{7\pi^2}{8 \cdot 30} g T^4. \quad (13)$$

So we have

$$\rho_b = \frac{8}{7} \rho_f = \frac{\pi^2}{30} g T^4, \quad (14)$$

where the scaling with the temperature again agrees with the simple dimensional analysis we performed in the last lecture.

Finally, for the pressure  $P$  we note that in the relativistic limit  $p^2/E(p) = p = E(p)$ , so that it trivially follows from the definitions in equations (3) and (4) that for bosons as well as fermions

$$P = \frac{1}{3} \rho = w \rho, \quad (15)$$

which nicely agrees with the equation of state parameter  $w = \frac{1}{3}$  for radiation.

## 1.2 Non-relativist particles

We can also analytically solve for  $n$ ,  $\rho$  and  $P$  in the non-relativistic limit, i.e. for regular matter. In this case we have

$$E(p) = \sqrt{m^2 + p^2} \approx m + \frac{p^2}{2m}. \quad (16)$$

Let us define  $x = p/\sqrt{2mT}$ . Since the temperature is related to the average kinetic energy  $T \sim \frac{p_{av}^2}{2m}$ , which is much smaller than  $m$ , we find that  $e^{E/T} \approx e^{m/T} \gg 1$ . This means that the distribution function, as given in (1), is the same for bosons and fermions

$$f(p) = \frac{1}{e^{E(p)/T} \pm 1} \approx e^{-\frac{E}{T}} \approx e^{-\frac{m}{T}} e^{-x^2}. \quad (17)$$

This then gives for the number density

$$n = \frac{g}{(2\pi)^3} e^{-\frac{m}{T}} \int_0^\infty 4\pi(2mT)^{\frac{3}{2}} x^2 e^{-x^2} dx = \frac{ge^{-\frac{m}{T}} (2mT)^{\frac{3}{2}} \Gamma\left(\frac{3}{2}\right)}{4\pi^2} = g \left(\frac{mT}{2\pi}\right)^{\frac{3}{2}} e^{-\frac{m}{T}}, \quad (18)$$

where we used that  $\Gamma(3/2) = \sqrt{\pi}/2$ .

In order to calculate the energy density, we use that  $E \approx m$  and find to leading order from the definition in equation (3) that

$$\rho = mn = gm \left(\frac{mT}{2\pi}\right)^{\frac{3}{2}} e^{-\frac{m}{T}}. \quad (19)$$

Finally, we again calculate the pressure  $P$  as given in (4). Here we use that  $p^2/E \approx p^2/m = (2mT)x^2/m$  and find, using the simplification leading to (18), that

$$P = \frac{g}{(2\pi)^3} \frac{e^{-\frac{m}{T}}}{3m} \int_0^\infty 4\pi(2mT)^{\frac{5}{2}} x^4 e^{-x^2} dx = \frac{ge^{-\frac{m}{T}} (2mT)^{\frac{5}{2}} \Gamma\left(\frac{5}{2}\right)}{12\pi^2 m} = gT \left(\frac{mT}{2\pi}\right)^{\frac{3}{2}} e^{-\frac{m}{T}} = nT, \quad (20)$$

where we used that  $\Gamma(5/2) = 3\sqrt{\pi}/4$ . Note that this is the familiar ideal gas law  $P = nk_B T$  or after multiplying by the volume  $V$ :  $PV = Nk_B T$ .

Now as we argued above, the temperature  $T$  is much smaller than the mass  $m$ . This means that

$$P = nT = \frac{T}{m} \rho = w\rho \approx 0 \quad \text{for } T \ll m. \quad (21)$$

So we again reproduce our previous result that for non-relativistic matter we can neglect the pressure.

The exponential decay of the number density for non-relativistic particles that we found in (18) can be understood as particle-anti-particle annihilation. As mentioned above, at high energies particle-anti-particle pairs are created and destroyed at equal rates but once the universe cools so much that the particle's mass starts to become important the pair creation stops and the annihilation leads to an exponential suppression in the number density of the particle.

## 2 The effective number of relativistic species

The total radiation density is given by the sum over the contributions from all particles

$$\rho_r = \sum_i \rho_i = \frac{\pi^2}{30} g_\star(T) T^4, \quad (22)$$

where  $i$  runs over all standard model particles and  $g_\star(T)$  is the effective number of degrees of freedom at temperature  $T$ , which we take to be the photon temperature. The sum over  $i$  can receive two contributions. One from relativistic particles that are in equilibrium with the photons, i.e. that have  $T_i = T \gg m_i$ . These contribute to  $g_\star$  the following

$$g_\star^{th}(T) = \sum_{i=bosons} g_i + \frac{7}{8} \sum_{i=fermions} g_i = g_b + g_f, \quad (23)$$

where *th* stand for thermal equilibrium. However, particles can decouple so that they won't be in thermal equilibrium with the photons anymore. If these particles are relativistic, i.e. we have  $T_i \neq T$  and  $T_i \gg m_i$ , then they contribute to  $g_\star$

$$g_\star^{dec}(T) = \sum_{i=bosons} g_i \left(\frac{T_i}{T}\right)^4 + \frac{7}{8} \sum_{i=fermions} g_i \left(\frac{T_i}{T}\right)^4, \quad (24)$$

where *dec* stands for decoupled. We thus have

$$g_\star(T) = g_\star^{th}(T) + g_\star^{dec}(T). \quad (25)$$

As we discussed last time at  $T \gg 100\text{GeV}$  all standard model particles are relativistic (cf. figure 3) and in thermal equilibrium with the photons (and each other).

So let us calculate  $g_\star$  for the standard model. We have the following contribution to  $g_b$ :

- The  $Z$ ,  $W^+$ ,  $W^-$  and the photon  $\gamma$  are all massless vectors and have two degrees of freedom each. Therefore they contribute  $4 * 2 = 8$ .
- Before the electroweak phase transition the Higgs scalar is a two vector whose entries are complex scalars so that it contributes  $2 * 2 = 4$ .
- There are actually 8 gluons<sup>1</sup> that are all massless vectors, so that they contribute  $8 * 2 = 16$ .

This leads to a total of  $g_b = 28$ .

Massive fermions have two possible spins and therefore have each two internal degrees of freedom each. We take the neutrinos to be only left-handed, i.e. we assume that the right-handed neutrinos are very heavy and do not contribute. Fermions also have antiparticles that we need to include in our counting. Then we find the following contributions to  $g_f$ :

- The left-handed neutrinos and their anti-particles contribute  $3 * 2 = 6$ .

---

<sup>1</sup>This is not clear from figure 3 and follows from the more complicated nature of the strong force. The gluons are the entries in an  $SU(3)$  matrix that has eight real independent entries.




Figure 1: The known particles in our universe (taken from Wikipedia).

- The electron  $e$ , the  $\mu$  and the  $\tau$  contribute twice this  $3 * 2 * 2 = 12$ .
- The six quarks can have three distinct charges under the strong force<sup>2</sup> which leads to an additional factor of 3 so that we have  $6 * 2 * 2 * 3 = 72$ .

Thus in total we have  $g_f = 90$  and the value of  $g_*$  at temperatures well above a  $100\text{GeV}$  is

$$g_* = g_b + \frac{7/8}{g_f} = 28 + \frac{7}{8}90 = 106.75. \quad (26)$$

In an expanding and cooling universe particles will become non-relativistic. Before we discuss this in detail in the next subsection, let us mention the electroweak phase transition: At a temperature around  $100\text{GeV}$  the standard model of particle physics undergoes a transition during which the Higgs field develops a vacuum expectation value. This vacuum expectation value is actually what gives a mass to all the fields (particles) in the standard model. After this phase transition the  $W^\pm$  and  $Z$  gauge fields have a mass. Massive vectors have three internal degrees of freedom so that this modifies our counting above. However, these new three degrees of freedom (one for each  $W^+$ ,  $W^-$  and  $Z$ ) come from the Higgs field that after this transition is only left with a single degree of freedom. So the net number of degrees of freedom does not change during the electroweak phase transition.

<sup>2</sup>Each of them is a three vectors on which the gluon  $SU(3)$  matrix can act.

### 3 Particle freeze-out

Once the temperature of the universe drops below the mass of a particle, the particle-anti-particle annihilation for this particle is favored compared to particle-anti-particle creation. This leads to an exponential decay of the particle number  $n$ , as derived in equation (18). This transition from relativistic to non-relativistic particle and the resulting annihilation of particles with their anti-particles is not instantaneous. Roughly 80% of the particles are annihilated in the interval  $m > T > m/6$ .

One effect of this so called particle freeze out is, as we will discuss next time, that the decrease in temperature of the universe is slowed down, since the particle-anti-particle annihilation deposits the energy contained in the annihilating particles into the remaining particles that are still in thermal equilibrium. But what happens to the particles themselves? Do they completely disappear?

In a non-expanding (but still somehow cooling) universe with vanishing chemical potential for these particles, this would be the case and the number density would keep decreasing exponentially with the temperature. However, as we mentioned last time (cf. equation (12) in the lecture 5 notes), in our universe there is a small baryon-anti-baryon asymmetry so that the particles cannot all annihilate since there aren't enough anti-particles around. This leads to the observed remaining baryons in our universe. Note, that in the standard model of particle physics heavy quarks (and leptons) decay to the lighter quarks (and leptons). For example, the top quark has an estimated lifetime of  $5 \times 10^{-25} s$  so that any relic top quarks will quickly decay to up quarks.

Another fate of non-relativistic particles in an expanding universe is that at a certain point their interaction rate  $\Gamma$  (which is proportional to their exponentially decaying number density  $n$ ) becomes so small that it is smaller than the Hubble expansion  $H$ . In such a case the particles and anti-particles cannot find each other anymore and the annihilation stops. The exponential decay in the particle density followed by this so called freeze-out is shown in the log-log-plot in figure 2.




Figure 2: Once a particle becomes non-relativistic its number density decays exponentially. The Hubble expansion or a non-zero chemical potential can lead to a relic density.

## 4 Evolution of the relativistic degrees of freedom

Having briefly discussed the potential fate of relativistic matter that becomes non-relativistic, let us return to the relativistic degrees of freedom of the standard model, during the time when our universe cools from a few hundred  $GeV$  to a few  $eV$ . The behavior of  $g_*(T)$  is shown in figure 3.




Figure 3: The evolution of the relativistic degrees of freedom in our early universe (taken from Daniel Baumann’s “Cosmology” lectures).

After the electroweak phase transition particles have their usual mass and the heaviest field, the top quark starts to become non-relativistic. This reduces  $g_*$  by  $12 * 7/8 = 10.5$ . Next the massive vector bosons  $W^\pm$  and  $Z$  and the Higgs scalar become non-relativistic, which reduces  $g_*$  by another  $9 + 1 = 10$ . After that the  $b$  and  $c$  quarks and the  $\tau$  become also non-relativistic. At a temperature of roughly  $150 MeV$  our universe undergoes another phase transition. The strong force becomes so strong that all quarks and gluons combine into uncharged bound states. For example, the  $u$  and  $d$  quarks combine into protons  $uud$  and neutrons  $udd$ . All the particles that are combinations of three quarks are called baryons and even the lightest of them, the proton, has a mass of  $1 GeV \gg 150 MeV$  so that after the QCD phase transition the baryons are all non-relativistic. However, there are also bound states of quarks and anti-quarks that are called mesons. The lightest mesons are the pi-mesons  $\pi^+ = u\bar{d}$ ,  $\pi^- = d\bar{u}$  and the  $\pi^0$  which is a combination of  $u\bar{u}$  and  $d\bar{d}$ . These three mesons have a mass of  $135 MeV - 140 MeV$  so that they will still be relativistic after the QCD transition. They become non-relativistic shortly before the  $\mu$  leaving only the electron  $e$ , the photons and neutrinos as relativistic particles. As you can see from the graph something interesting is happening once the electrons become non-relativistic and we will discuss this next time.