

Name:

DUE ON 29.4.2019

Matrikelnummer:

Homework - 5

1. The largest particle accelerator in the world, the Large Hadron Collider at CERN can reach energies of roughly 10TeV . So we understand and can experimentally test physics up to this energy scale. This means in turn that we essentially understand the evolution of our universe up to this energy scale very well.

The matter-radiation equality happened at redshift $z = 3400$, which corresponds to $t_{eq} = 6 \times 10^4$ years after the big bang. Before that our universe was radiation dominated $a(t) \approx a_{eq} \sqrt{t/t_{eq}}$ for $t < t_{eq}$. Determine the time *in seconds* at which $T_{CMB} = 10\text{TeV}$.

2. The early universe was transitioning from a radiation to a matter dominated phase. Let us derive an analytic solution for the scale factor for such a two component universe.

- (a) Rewrite the Friedmann equations (given in equations (1) and (2) in the lecture 2 notes) using conformal time.

- (b) Now assume only radiation and matter as source terms (i.e. set $K = 0$). Check that the second equation involving $a''(\tau)$ is not sourced by the radiation but only couples to matter. Simplify this equation using that $\rho_m a(\tau)^3 = \text{const.} = \frac{1}{2} \rho_{eq} a_{eq}^3$, where the subscript eq refers to the time of matter and radiation equality.

- (c) Solve this trivial differential equation. Fix one of the two integration constants by demanding that $a(\tau = 0) = 0$.

- (d) Fix the second integration constant by using the first Friedmann equation and the fact that

$$\rho = \rho_m + \rho_{rad} = \frac{\rho_{eq}}{2} \left[\left(\frac{a_{eq}}{a} \right)^3 + \left(\frac{a_{eq}}{a} \right)^4 \right]. \quad (1)$$

Express your answer in terms of a_{eq} and τ/τ_* with

$$\tau_* \equiv \sqrt{\frac{3}{\pi G \rho_{eq} a_{eq}^2}} = \frac{\tau_{eq}}{\sqrt{2-1}}. \quad (2)$$

3. During the early universe the standard model of particle physics undergoes phase transitions. In particular during the so called electroweak phase transition, the Higgs fields rolls down its potential and thereby reduces the cosmological constant from $\lambda_{before} \approx 10^{-65}$ to $\lambda_{today} \approx 10^{-120}$. This happens during the radiation dominated era of our universe at time t_{EW} when the temperature of the expanding universe drops below a critical temperature. Assume $\rho_c(t_{EW}) \sim \rho_{rad}(t_{EW}) \sim \rho_\Lambda(t_{EW})$ and explain in words, why such a transition does not affect the evolution of our universe at neither earlier nor later times. Redraw figure 2 in the lecture notes of lecture 4 including such a phase transition in the early universe.